

Inflahil

Ibuprofen

Composition

Inflahil 200 Tablet: Each film coated tablet contains Ibuprofen USP 200 mg.

Inflahil 400 Tablet: Each film coated tablet contains Ibuprofen USP 400 mg.

Inflahil Suspension: Each 5ml suspension contains Ibuprofen USP 100 mg.

Indications and Uses

Inflahil is used for its analgesic and anti-inflammatory effect in the treatment of rheumatoid arthritis, juvenile rheumatoid arthritis, ankylosing spondylitis, osteoarthritis and other non-rheumatic conditions, such as frozen shoulder (Capsulitis), bursitis, tendonitis, tenosynovitis and low back pain. Inflahil is indicated in the relief of mild to moderate pain such as dysmenorrhoea, dental and post-operative pain and in the relief of migraine.

Dosage and Administration

Adults: The recommended dosage of Inflahil is 1200 mg to 1800 mg daily in divided doses. Some patients can be maintained on a daily dose of 600 to 1200 mg. In severe or acute conditions the dosage may be increased until the acute phase is brought under control. However, the total daily dosage should not exceed 2400 mg in divided doses.

Children: The daily dosage of Inflahil is 20 mg per kg of body-weight in divided doses. In juvenile rheumatoid arthritis upto 40 mg per kg of body-weight daily in divided doses may be given. In children weighing less than 30 kg the total daily dose should not exceed 500 mg.

Elderly: No special dosage modification is required for elderly patients unless there is an impairment of renal or hepatic function. In case of such impairment, the dosage should be assessed individually by the physician.

Use in pregnancy & lactation

Although no teratogenic effects have been demonstrated in animal experiments, the use of Ibuprofen during pregnancy should, if possible, be avoided. In the limited studies so far available, Ibuprofen appears in the breast milk in very low concentration and is unlikely to affect the breast-fed infant adversely.

Contraindications

Ibuprofen is contra-indicated in patients with a history of peptic ulceration. Bronchospasm may be precipitated in patients suffering from or with a previous history of bronchial asthma. Ibuprofen should not be given to patients in whom aspirin and other non-steroidal anti-inflammatory drugs induce the symptoms of asthma, rhinitis or urticaria.

Side-effects

The side-effects of Ibuprofen may include dyspepsia, gastro-intestinal intolerance and bleeding and skin rashes of various types. Less frequently, thrombocytopenia has occurred.

Overdosage

In case of overdosage, gastric lavage and if necessary, correction of serum electrolytes have been recommended. There is no specific antidote for Ibuprofen.

Drug interactions

In therapeutic doses there is no evidence of significant interactions with other commonly used drugs. However, caution should be exercised in patients receiving oral anticoagulants and thiazide diuretics.

Storage

Store below 30° C in a dry place. Keep away from light.

Warning

To be used as directed by the physician. Keep out of reach of children.

Packaging

Inflahil 200 Tablet: Each box contains 100 tablets in alu-pvc blister pack.

Inflahil 400 Tablet: Each box contains 100 tablets in alu-pvc blister pack.

Inflahil Suspension: Each box contains 100 ml suspension in a PET bottle and a measuring cup.



Manufactured by:

Goodman Pharmaceuticals Ltd.

Better Health For All
Bhangnahati, Sreepur, Gazipur, Bangladesh